



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2016

---

## **The Role of Molecular Diagnostics in the Management of Patients with Gliomas**

Wirsching, H G ; Weller, M

**Abstract:** The revised World Health Organization (WHO) classification of tumors of the central nervous system of 2016 combines biology-driven molecular marker diagnostics with classical histological cancer diagnosis. Reclassification of gliomas by molecular similarity beyond histological boundaries improves outcome prediction and will increasingly guide treatment decisions. This change in paradigms implies more personalized and eventually more efficient therapeutic approaches, but the era of molecular targeted therapies for gliomas is yet at its onset. Promising results of molecularly targeted therapies in genetically less complex gliomas with circumscribed growth such as subependymal giant cell astrocytoma or pilocytic astrocytoma support further development of molecularly targeted therapies. In diffuse gliomas, several molecular markers that predict benefit from alkylating agent chemotherapy have been identified in recent years. For example, co-deletion of chromosome arms 1p and 19q predicts benefit from poly-chemotherapy with procarbazine, CCNU (lomustine), and vincristine (PCV) in patients with anaplastic oligodendroglioma, and the presence of 1p/19q co-deletion was integrated as a defining feature of oligodendroglial tumors in the revised WHO classification. However, the tremendous increase in knowledge of molecular drivers of diffuse gliomas on genomic, epigenetic, and gene expression levels has not yet translated into effective molecular targeted therapies. Multiple reasons account for the failure of early clinical trials of molecularly targeted therapies in diffuse gliomas, including the lack of molecular entry controls as well as pharmacokinetic and pharmacodynamics issues, but the key challenge of specifically targeting the molecular backbone of diffuse gliomas is probably extensive clonal heterogeneity. A more profound understanding of clonal selection, alternative activation of oncogenic signaling pathways, and genomic instability is warranted to identify effective combination treatments and ultimately improve survival.

DOI: <https://doi.org/10.1007/s11864-016-0430-4>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126712>

Journal Article

Accepted Version

Originally published at:

Wirsching, H G; Weller, M (2016). The Role of Molecular Diagnostics in the Management of Patients with Gliomas. *Current Treatment Options in Oncology*, 17(10):51.

DOI: <https://doi.org/10.1007/s11864-016-0430-4>

## **The Role of Molecular Diagnostics in the Management of Patients with Gliomas**

Hans-Georg Wirsching, Michael Weller\*

Dr. Hans-Georg Wirsching  
Resident Physician  
Department of Neurology  
University Hospital Zurich  
Frauenklinikstrasse 26  
CH-8091 Zurich  
Switzerland  
Phone: + 41 44 255 5500  
Fax + 41 44 255 4507  
Email [hans-georg.wirsching@usz.ch](mailto:hans-georg.wirsching@usz.ch)

Prof. Dr. Michael Weller  
Chairman  
Department of Neurology  
University Hospital Zurich  
Frauenklinikstrasse 26  
CH-8091 Zurich  
Switzerland  
Phone: + 41 44 255 5500  
Fax + 41 44 255 4507  
Email [michael.weller@usz.ch](mailto:michael.weller@usz.ch)

\* corresponding author

Illustrating material:

Table 1 – Gliomas in the 2016 WHO classification of primary tumors of the central nervous system

Figure 1 – Overview of molecular subtypes of diffuse gliomas

## **Opinion statement**

The revised World Health Organization (WHO) classification of tumors of the central nervous system of 2016 combines biology driven molecular marker diagnostics with classical histological cancer diagnosis. Reclassification of gliomas by molecular similarity beyond histological boundaries improves outcome prediction and will increasingly guide treatment decisions. This change in paradigms implies more personalized and eventually more efficient therapeutic approaches, but the era of molecular targeted therapies for gliomas is yet at its onset. Promising results of molecularly targeted therapies in genetically less complex gliomas with circumscribed growth such as subependymal giant cell astrocytoma or pilocytic astrocytoma support further development of molecularly targeted therapies. In diffuse gliomas, several molecular markers that predict benefit from alkylating agent chemotherapy have been identified in recent years. For example, co-deletion of chromosome arms 1p and 19q predicts benefit from polychemotherapy with procarbazine, CCNU (lomustine) and vincristine (PCV) in patients with anaplastic oligodendroglioma, and the presence of 1p/19q co-deletion was integrated as a defining feature of oligodendroglial tumors in the revised WHO classification. However, the tremendous increase in knowledge of molecular drivers of diffuse gliomas on genomic, epigenetic and gene expression levels has not yet translated into effective molecular targeted therapies. Multiple reasons account for the failure of early clinical trials of molecularly targeted therapies in diffuse gliomas, including the lack of molecular entry controls as well as pharmacokinetic and pharmacodynamics issues, but the key challenge of specifically targeting the molecular backbone of diffuse gliomas is probably extensive clonal heterogeneity. A more profound understanding of clonal selection, alternative activation of oncogenic signaling pathways and genomic instability is warranted to identify effective combination treatments and ultimately improve survival.

## Introduction

Gliomas are a histologically and molecularly heterogeneous group of tumors that are thought to derive from neuroglial progenitor cells and that account for approximately 27% of all primary brain tumors and for 80% of malignant primary brain tumors [1]. Histologically, the WHO classification of tumors of the central nervous system distinguishes astrocytomas, oligodendrogliomas and ependymomas, and assigns WHO grades I-IV with respect to the grade of malignancy [2, 3]. By convention, WHO grade I and II gliomas were referred to as low-grade and grades III and IV as high-grade gliomas. Low-grade gliomas comprised all gliomas with circumscribed growth, and these are rare and molecularly less complex than diffuse gliomas. The vast majority of gliomas are, however, characterized by diffuse growth and heterogeneous molecular patterns, and tremendous progress of molecular profiling array technologies has yielded genomic, epigenetic and gene expression landscapes that have refined the understanding and classification of biologically distinct glioma subtypes beyond histology [4-6] (Figure 1). These advances were acknowledged with the 2016 revision of the WHO classification of tumors of the central nervous system, which includes novel classes of diffuse gliomas based on genomic features (Table 1). Molecular diagnostics trump discordant histological results and thereby increase diagnostic accuracy and prognostic yield compared to previous histology based classifications [3]. Accordingly, the division into low grade and high grade gliomas is outdated and should be abandoned.

In parallel to these advances in classifying gliomas, the identification of key drivers of malignancy led to novel therapeutic approaches to target the molecular machinery of gliomas. Success of such targeted therapies in the molecularly less complex gliomas with circumscribed growth support this principle, but the extensive intratumoral heterogeneity of diffuse gliomas poses major challenges: First, large numbers of

molecular alterations in each tumor complicate the differentiation between passenger and driver events. Second, targeting a single driver alteration may be ineffective due to compensating pathways that are active in parallel or can be readily activated. Third, successful targeting of a tumor cell population may be compensated by clonal selection of another population that does not bear the targeted alteration. Therefore, a profound understanding of the molecular events that initiate and drive the progression of gliomas is crucial for the development of molecularly targeted therapies, because ultimately only targeting of key disease drivers that are shared by all cells within the tumor, or combination therapies can overcome treatment resistance.

Innovative trial designs have been suggested, such as umbrella trials that incorporate molecular profiling at study entry and the subsequent choice of an individually tailored therapy that matches the molecular profile of the tumor [7]. The scope of such trial designs is exploratory, thus underscoring the necessity to include repeat biopsies to enable the assessment of mechanisms of escape or resistance to molecular targeted therapies. However, the choice of tested molecular alterations in such trial designs is subject to expert opinions and availability of valid inhibitors that, amongst others, match the pharmacokinetic issues posed by the blood brain barrier. Technological advances such as deep sequencing and single-cell array approaches in tissues derived from sequential biopsies or surgeries will shed further light into the mechanisms underlying therapy resistance and malignant progression, albeit the prospective applicability of such large-scale arrays for clinical decision making or as molecular entry controls for clinical trials is still limited by logistic issues such as the requirement of extensive financial, human and technological resources, as well as a lack of consensus on data interpretation.

This review will summarize the current knowledge of the molecular background of gliomagenesis, the applicability of molecular marker diagnostics derived from this knowledge as well as current and future implications for precision medicine in gliomas.

## Gliomas with circumscribed growth

Improved understanding of the molecular basis of gliomas with circumscribed growth yielded promising results of molecularly targeted therapies. For example, subependymal giant cell astrocytomas (SEGA) are driven by mutations in either of two tumor suppressor genes, tuberous sclerosis complex (*TSC*)-1 and -2 that encode hamartin and tuberin, respectively. Loss of either protein yields disinhibition of cell growth signaling through mammalian target of rapamycin (mTOR) [8], and the mTOR inhibitor everolimus is effective against SEGA [9, 10].

Pilocytic astrocytoma is likewise considered a single pathway disease that is driven by the mitogen-activated protein kinase (MAPK) pathway due to mutations in the fibroblast growth factor receptor 1 (*FGFR1*) or neurotrophic receptor tyrosine kinase 2 (*NTRK2*) genes, or due to tandem duplications in a v-Raf murine sarcoma viral oncogene homolog B (*BRAF*) fusion gene [11, 12] and an early phase study of the MAPK inhibitor selumetinib in pilocytic astrocytoma has recently completed accrual (NCT01089101).

Pleomorphic xanthoastrocytomas are rare and less well characterized compared to other glioma entities. Particularly the mechanisms underlying the rare instances of malignant progression remain unclear. However, activating *BRAF*<sup>V600E</sup> mutations are present in more than half of pleomorphic xanthoastrocytomas [13] and recent reports of response to vemurafenib in heavily pretreated patients with recurrent pleomorphic xanthoastrocytomas suggest that the presence of the *BRAF*<sup>V600E</sup> mutation lends a rationale for salvage treatment with drugs like vemurafenib [14, 15], but no prospective trials are ongoing to further explore the role of *BRAF*<sup>V600E</sup> as a putative molecular target in pleomorphic xanthoastrocytomas.

## Ependymomas

Low-grade ependymomas are characterized by circumscribed growth patterns, but diffuse growth and multifocal progression occur frequently in the course of disease. Assessment of the v-rel avian reticuloendotheliosis viral oncogene homolog A (*RELA*)–*C11orf95* fusion gene has recently entered clinical routine, because *RELA* fusion-positive ependymomas have been included in the revised WHO classification as an entity with high risk for rapid and diffuse growth at diagnosis or during the disease course [3, 16]. A total of 9 molecular ependymoma subtypes have been defined, including a second high-risk subtype designated posterior fossa A (PF-A). PF-A ependymomas comprise the vast majority of posterior fossa ependymomas, but a driver alteration of this probably epigenetically driven disease has not yet been identified [17]. The standard of care in ependymomas comprises maximum safe resection and post-operative radiotherapy [18, 19]. In consideration that approximately half of all patients with ependymomas are children and that maximum resection may indeed be curative in a substantial fraction of patients, the timing of radiotherapy is a key aspect of the management of ependymomas, but probably radiotherapy should not be delayed in PF-A and *RELA* fusion-positive high-risk patients. The role of chemotherapy for the treatment of ependymomas has remained elusive and mostly confined to the recurrent setting [19]. Ongoing trials assess carboplatin-based regimens with and without bevacizumab (NCT01295944, NCT01088035), everolimus (NCT02155920) as well as a combination of temozolomide and the dual epithelial growth factor receptor (EGFR) inhibitor lapatinib (NCT00826241), which is directed against the EGFR isotypes Erb1 and Erb2 (HER2/neu), but molecularly targeted therapies taking the recent definitions of ependymoma subtypes into account have not yet been launched. Molecularly guided trials are complicated by the overall low frequency of ependymomas and further restrictions implicated by the apparent molecular diversity.

## **Diffuse oligodendroglial and astrocytic gliomas**

Joint efforts of the scientific community in characterizing the complex biology of diffuse gliomas culminated in The Cancer Genome Atlas Research Network (TCGA) and recent integration of information on exome sequences, DNA copy numbers, epigenetics, as well as messenger RNA, microRNA and protein expression identified prognostic subgroups of diffuse gliomas that were defined more accurately than by histology [6, 20, 21] (Figure 1). Molecular markers that represent a surrogate excerpt of these complex analyses are warranted to enable the translation of knowledge derived from these complex large-scale analyses into clinical practice (Table 1). Most current studies come up with a simplified approach of characterizing three major classes of gliomas, defined by isocitrate dehydrogenase (IDH) mutation plus 1p/19q codeletion, IDH mutation without 1p/19q codeletion, or IDH wildtype status [20, 22-25]. A classification of 5 prognostic adult glioma subgroups has been proposed based on the presence of *IDH* mutations, 1p/19q co-deletions and telomerase reverse transcriptase (*TERT*) promoter mutations [26], and independently of this classification, *MGMT* promoter methylation has emerged as a predictive marker for benefit from temozolomide chemotherapy in elderly patients with glioblastoma [27, 28].

***IDH* mutations.** The integration of *IDH* mutations into the revised WHO classification as a primary categorizing molecular alteration of diffuse gliomas has underpinned a change in paradigms from a histology-based towards a biology-based categorization of gliomas [3]. The conceptual novelty of the 2016 WHO classification is exemplified by the close and causal association of mutant IDH with a distinct epigenetic pattern designated glioma-CpG island methylator phenotype (G-CIMP) [29, 30]. Oncogenesis studies identified *IDH* mutations as early and possibly initiating events during



gliomagenesis and demonstrated that the G-CIMP pattern is preserved throughout malignant progression of *IDH* mutated gliomas from WHO grades II through IV [5, 31]. The vast majority of histological WHO grade II/III gliomas harbor *IDH* mutations, but only about 10% of glioblastomas [32, 33]. The histological appearance of *IDH* mutated versus wild-type glioblastomas is largely identical, but *IDH* mutated glioblastomas are associated with younger age at diagnosis and a better prognosis. An antibody to specifically detect the most common *IDH* mutation in gliomas *IDH1*<sup>R132H</sup> by immunohistochemistry is commercially available [34] and additional sequencing to detect other *IDH1/2* mutations is recommended in patients with WHO grade II or III gliomas or young adults with glioblastoma whose tumors show negative immunohistochemistry [35]. To date, the detection of *IDH* mutations has no direct therapeutic implications, but refines the prognostic accuracy of the WHO classification.

**Co-deletion of chromosome arms 1p and 19q.** According to the 2016 WHO classification, co-deletion of chromosome arms 1p and 19q is a defining feature of oligodendroglial tumors, whereas 1p/19q non-co-deleted tumors are classified as astrocytomas [3]. Notably, this molecular definition of neuroglial lineage association has outdated the diagnosis of a mixed oligoastrocytic entity, the histological diagnosis of which has been debated because of inter-observer variations and which, according to the revised WHO classification, may now only be diagnosed in patients that lack assessment of 1p/19q status [3, 36]. 1p/19q co-deleted oligodendrogliomas constitute a sub-group of *IDH* mutated gliomas that commonly harbor additional activating mutations in the promoter region of *TERT*. In contrast, oncogenic mutations in the tumor suppressor gene *TP53* and in the methylome organizer alpha thalassemia/mental retardation syndrome X-linked (*ATRX*) are primarily found in *IDH* mutated astrocytomas and associated with less favorable prognosis [26, 36, 37]. Fluorescent *in situ* hybridization (FISH) and microsatellite analysis for loss of heterozygosity are commonly applied to assess 1p/19q status in clinical routine. The

prognosis of 1p/19q co-deleted gliomas is favorable and malignant progression to glioblastoma is rare. Besides diagnostic and prognostic implications, a predictive role of 1p/19q co-deletion for benefit from the addition of polychemotherapy with procarbazine, CCNU/lomustine and vincristine (PCV) to radiotherapy was identified in long-term analyses of two phase III trials in patients with anaplastic oligodendroglial tumors at a median follow-up time of more than ten years [38, 39]. Less toxic regimens were commonly utilized in clinical practice by the time these results have re-defined PCV as the standard of care for 1p/19q co-deleted tumors and the phase III CODEL trial is currently exploring whether PCV may be substituted by temozolomide to reduce toxicity, without compromising survival (NCT00887146).

**Mutations of the telomerase reverse transcriptase (*TERT*) promoter.** *TERT* promoter mutations are present in most diffuse gliomas and confer poor prognosis in patients with *IDH* wild-type, WHO grade II/III diffuse gliomas [26]. This is of particular relevance for the management of patients with WHO grade II diffuse gliomas, because watchful waiting strategies are considered feasible in patients with favorable prognostic features, including age < 40, tumor diameter < 6 cm, lack of midline crossing and lack of neurological deficit other than an brain tumor-associated epilepsy [40-42], but such strategies are probably not adequate in the poor prognosis subgroup of *TERT* mutated, *IDH* wild type diffuse gliomas.

**Hypermethylation of the O6-methylguanyl DNA methyltransferase (*MGMT*) promoter.** The DNA repair protein *MGMT* counteracts DNA damage conferred by alkylating agent chemotherapy. Transcriptional silencing of the *MGMT* gene due to promoter hypermethylation occurs almost invariably in G-CIMP positive, *IDH* mutated diffuse gliomas, as opposed to approximately 50% of G-CIMP negative, *IDH* wild-type diffuse gliomas [26, 43]. *MGMT* methylation status has a pivotal role for clinical decision-making, e.g., in elderly glioblastoma patients, but *MGMT* does not segregate

glioma subtypes with distinct gene ontologies. Consequently, *MGMT* methylation status was not included as a segregating feature in the revised WHO classification. Of note, different technical approaches and a lack of consensus on data interpretation render *MGMT* status assessments prone to variability [44]. *MGMT* promoter methylation was strongly associated with benefit from chemotherapy with temozolomide in the trial carried out by the European Organisation for Research and Treatment of Cancer (EORTC) and the National Cancer Institute of Canada (NCIC) that defined adjuvant temozolomide as the standard of care for newly diagnosed glioblastoma [45, 46]. In the elderly setting, the superiority of the combination of temozolomide and radiotherapy over radiotherapy alone has been confirmed [47]. Elderly patients with diffuse gliomas not considered candidates for combined modality treatment are commonly treated with monotherapy regimens [48]. Two phase III trials in elderly patients with mostly glioblastomas and a smaller fraction of anaplastic astrocytomas defined a predictive role of *MGMT* methylation for response to temozolomide versus radiotherapy to guide decision making at primary diagnosis [27, 28]. The strong association of *MGMT* methylation with benefit from temozolomide in glioblastoma is retained at recurrence, thus making rechallenge with temozolomide a viable option for patients with *MGMT* methylated glioblastoma and prolonged progression-free survival with first-line radiochemotherapy [49]. Changes of *MGMT* promoter methylation status at recurrence have not been observed in studies comparing matched tissue samples [50]. Interestingly, the prognostic value of *MGMT* methylation was similar among patients receiving radiotherapy versus chemotherapy in subgroup analyses of two phase III trials in patients with anaplastic gliomas [51, 52], albeit more global DNA methylation analyses suggest a predictive role of methylation of two specific CpG sites in the *MGMT* promoter for benefit from PCV [53]. However, the number of samples in this cohort was small and warrants confirmatory analyses. Overall, the role of *MGMT* promoter methylation status in WHO grade II and grade III gliomas remains a matter of controversy.

**Histone H3.3 (*H3F3A*) K27 and G34 mutations.** Point mutations in the 2 critical aminoacids K27 and G34 of the gene encoding histone H3.3 define *IDH* wild-type glioma subtypes in children and adolescents with G-CIMP negative gene methylation patterns that arise from distinct anatomical compartments [21, 54]. K27 and G34 mutations are mutually exclusive and strongly associated with mutations of the tumor suppressor gene *TP53* [21]. The strong anatomical and molecular association prompted the inclusion of *H3F3A*<sup>K27M</sup> mutant diffuse midline glioma as a novel entity to the 2016 WHO classification [3]. K27 mutations confer a poor prognosis, which may in part be attributed to the poorer surgical accessibility of midline tumors, but to date, no therapeutic implications derive from the presence of *H3F3A* mutations.

**Molecular subtypes derived from integrated large-scale analyses.** The complexity of diffuse gliomas is not fully captured by marker-based sub-classification. Integrated gene expression and genomic analyses defined 4 glioblastoma subtypes that were associated with distinct genomic alterations and that were termed neural, proneural, classical and mesenchymal based on similarities to previously defined developmental gene expression signatures [55]. Unsupervised cluster analyses of subsequent independent datasets did not reproduce the segregation of the neural subtype signature as a distinct entity [21, 56], but further integration of methylation data defined a total of 6 subtypes, including 4 that clustered with proneural gene expression and are more common in children or young adults [21]. *IDH* mutated glioblastomas were the only proneural glioblastomas to confer a favorable prognosis, whereas the prognostic less favorable subtypes with proneural gene expression were characterized by *H3F3A* mutations in the codons of the critical amino acids K27 or G34, or by amplification of the gene encoding platelet derived growth factor receptor A (*PDGFRA*). *PDGFRA* amplified glioblastomas were designated receptor tyrosine kinase (RTK) I, as opposed to the non-proneural RTK II subtype that is characterized

by a classical gene expression pattern and amplification of the receptor tyrosine kinase *EGFR*, and both RTK I and RTK II subtypes are furthermore associated with deletions of the cyclin-dependent kinase inhibitor 2A (*CDKN2A*) tumor suppressor gene and poor prognosis. The mesenchymal subtype shows a lower rate of copy-number alterations and no characteristic mutations, but likewise confers poor prognosis [21]. More recently, the applicability of these glioblastoma-based classifications to the entire histological spectrum of diffuse oligodendroglial and astrocytic diffuse gliomas was confirmed and two further prognostic subgroups were identified: (i) A small subgroup of *IDH* mutated astrocytic gliomas with poor prognosis that do not exhibit the characteristic G-CIMP gene methylation pattern, but are instead characterized by cell cycle activation due to *CDKN2A* deletions and cyclin dependent kinase 4 (*CDK4*) amplifications, and (ii) a small subgroup *IDH* wild-type WHO grade II/III astrocytic and oligodendroglial tumors in mostly young adults with favorable prognosis that share epigenetic and genomic features of pilocytic astrocytomas [6]. Such large-scale analyses rely on data derived from entire tissue samples, but another level of complexity is added by the extensive spatial and temporal molecular heterogeneity of gliomas [4, 5, 31].

### **Avenues and challenges of molecular targeted therapies in diffuse gliomas**

The perspective of specifically targeting disease drivers in diffuse gliomas evoked high hopes for future effective treatments, but to date such attempts have failed beyond early phase studies. The paramount reason for this failure is the molecular complexity and heterogeneity of diffuse gliomas, but a broad range of factors that can be overcome by careful planning of clinical trials have had their share, too. For example, pharmacokinetic and pharmacodynamics issues have complicated early trials of EGFR inhibition in glioblastoma: The glioma-specific *EGFRvIII* deletion

mutation that is present in half of all *EGFR* amplified glioblastomas yields constitutive, ligand-independent receptor activity that is not targeted by the EGFR inhibitors erlotinib or gefitinib, whereas brain penetration of the EGFRvIII inhibitor lapatinib is limited [57]. Another challenge in establishing molecular targeted therapies is posed by the choice of hypothesis-based or empirical molecular entry controls to choose patients that will potentially benefit from a trial drug. For example, the mammalian target of rapamycin (mTOR) inhibitor temsirolimus (CCI-779) failed to improve outcome in two separate phase II trial cohorts, whereas response was associated with phosphorylation of mTORSer2448 in newly diagnosed *MGMT* unmethylated glioblastoma [58] and with phosphorylation of the mTOR target S6-kinase in recurrent glioblastoma [59], and efficacy studies further exploring mTOR inhibition in glioblastoma patients pre-selected for these molecular markers are currently probably not underway.

Finally, choosing the right target poses the most difficult challenge to clinical trials that aim at specifically targeting the molecular machinery of cancers. In consideration of the extensive vascularity of glioblastomas, targeting signaling through the key molecular signaling molecule vascular endothelial growth factor (VEGF) was explored extensively, but a total of six phase III trials in newly diagnosed and recurrent glioblastoma failed to demonstrate an overall survival benefit from anti-angiogenic therapy targeting either the VEGF or other putative angiogenic pathways [60-65]. Molecular analyses have however identified the proneural gene expression signature as a potential molecular marker for the benefit of an adjunct of the anti-VEGF antibody bevacizumab to standard chemoradiotherapy in newly diagnosed glioblastoma [66], but a confirmatory trial that applies this signature prospectively is currently not planned and the opportunity to validate this finding in the RTOG 0825 trial is pending. Considering the prominent role of epigenetic mechanisms in the pathophysiology of diffuse gliomas, measures to induce epigenetic reshaping were successfully applied in preclinical studies, including a small molecule inhibitor of

mutant *IDH* [67] and an inhibitor of the K27 demethylase JMJD3, which yielded K27 hypermethylation and potent anti-tumor activity in a pre-clinical model of H3 K27M–mutant diffuse midline glioma [68], but clinical trials to test these inhibitors await to be completed.

Immunotherapy approaches have recently moved into the focus of novel developments in oncology: In patients with metastatic melanoma and non-small-cell lung cancer, monoclonal antibodies directed against the immunosuppressive T-cell receptors programmed death protein (PD)1 (nivolumab, pembrolizumab) or cytotoxic T-lymphocyte antigen (CTLA) 4 (ipilimumab) have proven a successful strategy to improve survival by promoting innate immune-mediated anti-tumor activity [69-73].

The efficacy of nivolumab is currently being assessed in newly diagnosed glioblastoma as a substitute for temozolomide in patients with an unmethylated *MGMT* promoter in the phase III trial CheckMate 498 (NCT02617589), and as an adjunct to temozolomide in patients with a methylated *MGMT* promoter in the phase II trial CheckMate 548 (NCT02667587). Furthermore, the phase III trial CheckMate 143 assessed the efficacy of nivolumab versus bevacizumab in patients with recurrent glioblastoma (NCT02017717).

Vaccination against tumor antigens is another immunotherapy approach that is being explored in glioblastoma. The double-blinded phase II ICT-107 trial explored the safety and efficacy of a vaccine consisting of autologous dendritic cells that were pulsed with six tumor-associated peptides to trigger an anti-tumor T-cell response in newly diagnosed glioblastoma. The adjunct of this vaccine to standard chemoradiotherapy prolonged progression-free survival by 2.4 months without relevant safety issues [74] and therefore led to a similar phase III trial (NCT02546102). The DCVax trial is a phase III trial that assesses the efficacy of vaccination with dendritic cells that were pulsed with patients' tumor lysates (NCT00045968). Using peptides for vaccination that occur specifically in tumor cells due to coding mutations potentially prevents cross reactivity of an elicited immune

response against normal host cells and renders immune tolerance less likely. However, due to the multi-branched ontology of diffuse gliomas, only few mutations that were acquired early during gliomagenesis are present in all clones [5], such as mutant IDH which could be targeted successfully in a pre-clinical glioma model [75]. In contrast, clonal selection is a likely mechanism of resistance to vaccines that aim to elicit immune responses against mutations acquired later during gliomagenesis, such as the *EGFRvIII* deletion. This limitation may underly the lack of efficacy of vaccination with the EGFRvIII-specific peptide rindopepimut as an adjunct to standard chemoradiotherapy in newly diagnosed EGFRvIII-positive glioblastoma (NCT01480479), albeit the same vaccine appeared to prolong survival as an adjunct to the anti-angiogenic and probably pro-immunogenic anti-VEGF antibody bevacizumab in the phase II ReACT trial in patients with recurrent glioblastoma [76]. The term adoptive T-cell transfer refers to an immunotherapy approach that exploits the *in vitro* expansion or genetic engineering of T-cells directed against tumor-associated or tumor-specific peptides, and chimeric antigen-receptor (CAR) T-cells are genetically engineered cells that furthermore circumvent the need for co-stimulatory molecules by linking antigen-binding domains directly to intracellular signaling domains that trigger T-cell activation [77]. A phase I trial in glioblastoma patients utilizing CAR T-cells directed against Her2 is ongoing (NCT01109095) and another trial utilizing CAR T-cells directed against EGFRvIII has suspended recruitment (NCT01454596).

In summary, the current quest for effective and feasible treatment modalities and targets is yet at the onset of the personalization of treatments in patients with diffuse gliomas. Probably only combination treatments guided by molecular fingerprints that predict or capture clonal heterogeneity will be able to overcome resistance, but a plethora of issues including logistic feasibility and toxicity will have to be resolved along this development.



**Figure 1. Overview of molecular subtypes of diffuse gliomas.** The layered classification included in the 2016 WHO classification is depicted in red. Diffuse gliomas are segregated by *IDH* mutation status. 1p/19q codeletion marks oligodendrogliomas and the *H3F3A*<sup>K27</sup> mutation marks diffuse midline gliomas. These molecular assessments trump histological tumor grades II-IV. Other less robust genetic and genomic (green), epigenetic (blue) and gene expression (yellow) patterns that have not been included in the WHO classification have been defined to further segregate diffuse gliomas based on molecular features. Relative prognosis and age associations are depicted at the bottom.

**Table 1. Gliomas in the 2016 WHO classification of primary tumors of the central nervous system [3].**

Entity	Histological WHO grade	Median overall survival	First line therapy	Comments
<b>Diffuse astrocytic and oligodendroglial tumours</b>				
Diffuse astrocytoma, IDH mutant	II	> 10 years [78]	Watchful waiting [40] or RT→PCV [79] or (TMZ)/RT→TMZ per extrapolation from WHO grade III gliomas [80]	The previous term fibrillary astrocytoma is discouraged, because fibrillary morphology is the classical appearance and not a variant.
Gemistocytic astrocytoma, IDH mutant	II	< 4 years [81]	RT→PCV [79] or (TMZ)/RT→TMZ per extrapolation from WHO grade III gliomas [80]	Variant of diffuse astrocytoma. Higher incidence of progression to IDH-mutant anaplastic astrocytoma and glioblastoma has been reported [81]. Survival times including IDH-mutation status have not been reported. Watchful waiting strategies may not be adequate, if less favorable prognosis is confirmed in IDH mutated gemistocytic astrocytoma.
Diffuse astrocytoma, IDH wild-type	II	> 5 years [79]	RT→PCV [79] or TMZ/RT→TMZ per extrapolation from IDH wild-type glioblastoma [46]	Rare, provisional entity comprising a variety of molecular subtypes mostly resembling IDH wild-type glioblastomas [20, 82].

				Watchful waiting strategies are probably not adequate. Survival times have not been reported.
Diffuse astrocytoma, NOS	II	4-5 years [1]	RT→PCV [79] or (TMZ)/RT→TMZ per extrapolation from WHO grade III gliomas [80]	
Anaplastic astrocytoma, IDH mutant	III	5-10 years [33, 78]	(TMZ)/RT→TMZ [80]	No population-based survival data reported.
Anaplastic astrocytoma, IDH wild-type	III	2-4 years [26, 33, 83]	TMZ/RT→TMZ per extrapolation from IDH wild-type glioblastoma [46], possibly RT alone in MGMT unmethylated patients [84]	Molecularly resembling IDH wild-type glioblastoma, <i>TERT</i> promoter mutations confer poor prognosis [20, 82]. No population-based survival data reported.
Anaplastic astrocytoma, NOS	III	1-2 years [1]	(TMZ)/RT→TMZ [80]	
Glioblastoma, IDH wild-type	IV	11-15 months [33, 85, 86]	TMZ/RT→TMZ [46, 47], > 65-70 years RT (MGMT unmethylated), or TMZ (MGMT methylated), if combination therapy is not deemed feasible [27, 28]	Histological variants: Giant cell glioblastoma, gliosarcoma, epitheloid glioblastoma
Glioblastoma, IDH mutant	IV	2-3 years [33, 85]	(TMZ)/RT→TMZ per extrapolation from IDH mutant anaplastic astrocytoma [80]	
Glioblastoma, NOS	IV	< 1 year [1, 86]	See above	
Diffuse midline glioma, H3-K27M mutant	IV	< 1 year [21]	TMZ/RT→TMZ per extrapolation from IDH wild-type glioblastoma [46, 47]	No population-based survival data reported.

Oligodendroglioma, IDH mutant and 1p/19q codeleted	II	> 10 years [79]	Watchful waiting [40] or RT→PCV [79]	No population-based survival data reported.
Oligodendroglioma, NOS	II	> 10 years [1, 87]	Watchful waiting [40] or RT→PCV [79]	
Anaplastic oligodendroglioma, IDH mutant and 1p/19q codeleted	III	> 10 years [38, 39, 83]	RT→PCV [38, 39]	No population-based survival data reported.
Anaplastic oligodendroglioma, NOS	III	5-7 years [1, 83]	RT→PCV [38, 39]	
Oligoastrocytoma, NOS	II	< 7 years [81]	Watchful waiting [40] or RT→PCV [79] per extrapolation from 1p/19q codeleted oligodendroglioma	Large interobserver variability of histological diagnosis [36, 88].
Anaplastic oligoastrocytoma, NOS	III	> 5 years [89]	RT→PCV [38, 39] per extrapolation from 1p/19q codeleted anaplastic oligodendroglioma	Large interobserver variability of histological diagnosis [36, 88] No population-based survival data reported.

NOS, not otherwise specified (Histological diagnosis in cases where molecular data have not been fully assessed)

## References

1. Ostrom QT, Gittleman H, Fulop J, Liu M, Blanda R, Kromer C, et al. CBTRUS Statistical Report: Primary Brain and Central Nervous System Tumors Diagnosed in the United States in 2008-2012. *Neuro-oncology*. 2015;17 Suppl 4:iv1-iv62.
  2. Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, et al. The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol*. 2007;114(2):97-109.
  3. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016;131(6):803-20.
- \*\* This reference summarizes the most important implications of the revised WHO classification of diffuse gliomas.
4. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, et al. Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science*. 2014;344(6190):1396-401.
  5. Bai H, Harmanici AS, Erson-Omay EZ, Li J, Coskun S, Simon M, et al. Integrated genomic characterization of IDH1-mutant glioma malignant progression. *Nat Genet*. 2016;48(1):59-66.

\* Patterns of the genomic evolution of IDH-mutant diffuse gliomas were identified in this study.

6. Ceccarelli M, Barthel FP, Malta TM, Sabedot TS, Salama SR, Murray BA, et al. Molecular Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma. *Cell*. 2016;164(3):550-63.

\* Two previously unrecognized molecular subtypes of diffuse gliomas were identified by integration of different level molecular data.

7. Galanis E, Wu W, Cloughesy T, Lamborn K, Mann B, Wen PY, et al. Phase 2 trial design in neuro-oncology revisited: a report from the RANO group. *Lancet Oncol*. 2012;13(5):e196-204.

8. Chan JA, Zhang H, Roberts PS, Jozwiak S, Wieslaw G, Lewin-Kowalik J, et al. Pathogenesis of tuberous sclerosis subependymal giant cell astrocytomas: biallelic inactivation of TSC1 or TSC2 leads to mTOR activation. *J Neuropathol Exp Neurol*. 2004;63(12):1236-42.

9. Krueger DA, Care MM, Holland K, Agricola K, Tudor C, Mangeshkar P, et al. Everolimus for subependymal giant-cell astrocytomas in tuberous sclerosis. *The New England journal of medicine*. 2010;363(19):1801-11.

10. Franz DN, Belousova E, Sparagana S, Bebin EM, Frost M, Kuperman R, et al. Everolimus for subependymal giant cell astrocytoma in patients with tuberous sclerosis complex: 2-year open-label extension of the randomised EXIST-1 study. *Lancet Oncol*. 2014;15(13):1513-20.

11. Jones DT, Hutter B, Jager N, Korshunov A, Kool M, Warnatz HJ, et al. Recurrent somatic alterations of FGFR1 and NTRK2 in pilocytic astrocytoma. *Nat Genet*. 2013;45(8):927-32.

12. Jones DT, Kocialkowski S, Liu L, Pearson DM, Backlund LM, Ichimura K, et al. Tandem duplication producing a novel oncogenic BRAF fusion gene

defines the majority of pilocytic astrocytomas. *Cancer research*.

2008;68(21):8673-7.

13. Schindler G, Capper D, Meyer J, Janzarik W, Omran H, Herold-Mende C, et al. Analysis of BRAF V600E mutation in 1,320 nervous system tumors reveals high mutation frequencies in pleomorphic xanthoastrocytoma, ganglioglioma and extra-cerebellar pilocytic astrocytoma. *Acta Neuropathol*. 2011;121(3):397-405.

14. Chamberlain MC. Salvage therapy with BRAF inhibitors for recurrent pleomorphic xanthoastrocytoma: a retrospective case series. *J Neurooncol*. 2013;114(2):237-40.

15. Lee EQ, Ruland S, LeBoeuf NR, Wen PY, Santagata S. Successful Treatment of a Progressive BRAF V600E-Mutated Anaplastic Pleomorphic Xanthoastrocytoma With Vemurafenib Monotherapy. *J Clin Oncol*. 2016;34(10):e87-9.

16. Parker M, Mohankumar KM, Punchihewa C, Weinlich R, Dalton JD, Li Y, et al. C11orf95-RELA fusions drive oncogenic NF-kappaB signalling in ependymoma. *Nature*. 2014;506(7489):451-5.

17. Pajtler KW, Witt H, Sill M, Jones DT, Hovestadt V, Kratochwil F, et al. Molecular Classification of Ependymal Tumors across All CNS Compartments, Histopathological Grades, and Age Groups. *Cancer Cell*. 2015;27(5):728-43.

\* Prognostic molecular subtypes of ependymomas were identified by integration of different level molecular data.

18. Merchant TE, Mulhern RK, Krasin MJ, Kun LE, Williams T, Li C, et al. Preliminary results from a phase II trial of conformal radiation therapy and evaluation of radiation-related CNS effects for pediatric patients with localized ependymoma. *J Clin Oncol.* 2004;22(15):3156-62.
19. Timmermann B, Kortmann RD, Kuhl J, Meisner C, Slavc I, Pietsch T, et al. Combined postoperative irradiation and chemotherapy for anaplastic ependymomas in childhood: results of the German prospective trials HIT 88/89 and HIT 91. *International journal of radiation oncology, biology, physics.* 2000;46(2):287-95.
20. Cancer Genome Atlas Research N, Brat DJ, Verhaak RG, Aldape KD, Yung WK, Salama SR, et al. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *The New England journal of medicine.* 2015;372(26):2481-98.
21. Sturm D, Witt H, Hovestadt V, Khuong-Quang DA, Jones DT, Konermann C, et al. Hotspot mutations in H3F3A and IDH1 define distinct epigenetic and biological subgroups of glioblastoma. *Cancer Cell.* 2012;22(4):425-37.
22. Wiestler B, Capper D, Sill M, Jones DT, Hovestadt V, Sturm D, et al. Integrated DNA methylation and copy-number profiling identify three clinically and biologically relevant groups of anaplastic glioma. *Acta Neuropathol.* 2014;128(4):561-71.
23. Suzuki H, Aoki K, Chiba K, Sato Y, Shiozawa Y, Shiraishi Y, et al. Mutational landscape and clonal architecture in grade II and III gliomas. *Nat Genet.* 2015;47(5):458-68.



24. Weller M, Weber RG, Willscher E, Riehmer V, Hentschel B, Kreuz M, et al. Molecular classification of diffuse cerebral WHO grade II/III gliomas using genome- and transcriptome-wide profiling improves stratification of prognostically distinct patient groups. *Acta Neuropathol.* 2015;129(5):679-93.
25. Weller M, Wick W, Aldape K, Brada M, Berger M, Pfister SM, et al. Glioma. *Nat Rev Dis Primers.* 2015;1:15017.
26. Eckel-Passow JE, Lachance DH, Molinaro AM, Walsh KM, Decker PA, Sicotte H, et al. Glioma Groups Based on 1p/19q, IDH, and TERT Promoter Mutations in Tumors. *The New England journal of medicine.* 2015;372(26):2499-508.
- \*\* This study provides a set of three molecular markers to define prognostic glioma subtypes.
27. Malmstrom A, Gronberg BH, Marosi C, Stupp R, Frappaz D, Schultz H, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol.* 2012;13(9):916-26.
28. Wick W, Platten M, Meisner C, Felsberg J, Tabatabai G, Simon M, et al. Temozolomide chemotherapy alone versus radiotherapy alone for malignant astrocytoma in the elderly: the NOA-08 randomised, phase 3 trial. *Lancet Oncol.* 2012;13(7):707-15.
29. Noushmehr H, Weisenberger DJ, Diefes K, Phillips HS, Pujara K, Berman BP, et al. Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell.* 2010;17(5):510-22.

30. Turcan S, Rohle D, Goenka A, Walsh LA, Fang F, Yilmaz E, et al. IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. *Nature*. 2012;483(7390):479-83.

31. Mazon T, Pankov A, Johnson BE, Hong C, Hamilton EG, Bell RJ, et al. DNA Methylation and Somatic Mutations Converge on the Cell Cycle and Define Similar Evolutionary Histories in Brain Tumors. *Cancer Cell*. 2015;28(3):307-17.

\* The extensive clonal heterogeneity of diffuse gliomas is elucidated by this study.

32. Parsons DW, Jones S, Zhang X, Lin JC, Leary RJ, Angenendt P, et al. An integrated genomic analysis of human glioblastoma multiforme. *Science*. 2008;321(5897):1807-12.

33. Yan H, Parsons DW, Jin G, McLendon R, Rasheed BA, Yuan W, et al. IDH1 and IDH2 mutations in gliomas. *The New England journal of medicine*. 2009;360(8):765-73.

34. Capper D, Zentgraf H, Balss J, Hartmann C, von Deimling A. Monoclonal antibody specific for IDH1 R132H mutation. *Acta Neuropathol*. 2009;118(5):599-601.

35. Felsberg J, Wolter M, Seul H, Friedensdorf B, Goppert M, Sabel MC, et al. Rapid and sensitive assessment of the IDH1 and IDH2 mutation status in cerebral gliomas based on DNA pyrosequencing. *Acta Neuropathol*. 2010;119(4):501-7.

36. Sahm F, Reuss D, Koelsche C, Capper D, Schittenhelm J, Heim S, et al. Farewell to oligoastrocytoma: in situ molecular genetics favor classification

as either oligodendroglioma or astrocytoma. *Acta Neuropathol.*

2014;128(4):551-9.

\* The requirement of oligoastrocytoma as a separate histological entity was omitted by this study.

37. Reuss DE, Sahm F, Schrimpf D, Wiestler B, Capper D, Koelsche C, et al. ATRX and IDH1-R132H immunohistochemistry with subsequent copy number analysis and IDH sequencing as a basis for an "integrated" diagnostic approach for adult astrocytoma, oligodendroglioma and glioblastoma. *Acta Neuropathol.* 2015;129(1):133-46.

38. van den Bent MJ, Brandes AA, Taphoorn MJ, Kros JM, Kouwenhoven MC, Delattre JY, et al. Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol.* 2013;31(3):344-50.

39. Cairncross G, Wang M, Shaw E, Jenkins R, Brachman D, Buckner J, et al. Phase III trial of chemoradiotherapy for anaplastic oligodendroglioma: long-term results of RTOG 9402. *J Clin Oncol.* 2013;31(3):337-43.

\*\* [38, 39] These studies defined the predictive value of 1p/19q codeletions for response to polychemotherapy.

40. van den Bent MJ, Afra D, de Witte O, Ben Hassel M, Schraub S, Hoang-Xuan K, et al. Long-term efficacy of early versus delayed radiotherapy for low-grade astrocytoma and oligodendroglioma in adults: the EORTC 22845 randomised trial. *Lancet.* 2005;366(9490):985-90.

41. Daniels TB, Brown PD, Felten SJ, Wu W, Buckner JC, Arusell RM, et al. Validation of EORTC prognostic factors for adults with low-grade glioma: a report using intergroup 86-72-51. *International journal of radiation oncology, biology, physics*. 2011;81(1):218-24.
42. Pignatti F, van den Bent M, Curran D, Debruyne C, Sylvester R, Therasse P, et al. Prognostic factors for survival in adult patients with cerebral low-grade glioma. *J Clin Oncol*. 2002;20(8):2076-84.
43. Bady P, Sciuscio D, Diserens AC, Bloch J, van den Bent MJ, Marosi C, et al. MGMT methylation analysis of glioblastoma on the Infinium methylation BeadChip identifies two distinct CpG regions associated with gene silencing and outcome, yielding a prediction model for comparisons across datasets, tumor grades, and CIMP-status. *Acta Neuropathol*. 2012;124(4):547-60.
44. Wick W, Weller M, van den Bent M, Sanson M, Weiler M, von Deimling A, et al. MGMT testing--the challenges for biomarker-based glioma treatment. *Nature reviews Neurology*. 2014;10(7):372-85.
45. Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *The New England journal of medicine*. 2005;352(10):997-1003.
46. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *The New England journal of medicine*. 2005;352(10):987-96.
47. Perry JR, Laperriere N, O'Callaghan CJ, Brandes AA, Menten J, Phillips C, et al. A phase III randomized controlled trial of short-course radiotherapy with or without concomitant and adjuvant temozolomide in

elderly patients with glioblastoma (CCTG CE.6, EORTC 26062-22061, TROG 08.02, NCT00482677). J Clin Oncol. 2016;34:suppl; abstr LBA2.

\* This study defines the efficacy of combined chemoradiotherapy in elderly patients with MGMT methylated and unmethylated glioblastoma.

48. Weller M, van den Bent M, Hopkins K, Tonn JC, Stupp R, Falini A, et al. EANO guideline for the diagnosis and treatment of anaplastic gliomas and glioblastoma. Lancet Oncol. 2014;15(9):e395-403.

49. Weller M, Tabatabai G, Kastner B, Felsberg J, Steinbach JP, Wick A, et al. MGMT Promoter Methylation Is a Strong Prognostic Biomarker for Benefit from Dose-Intensified Temozolomide Rechallenge in Progressive Glioblastoma: The DIRECTOR Trial. Clinical cancer research : an official journal of the American Association for Cancer Research. 2015;21(9):2057-64.

50. Felsberg J, Thon N, Eigenbrod S, Hentschel B, Sabel MC, Westphal M, et al. Promoter methylation and expression of MGMT and the DNA mismatch repair genes MLH1, MSH2, MSH6 and PMS2 in paired primary and recurrent glioblastomas. Int J Cancer. 2011;129(3):659-70.

51. Wick W, Hartmann C, Engel C, Stoffels M, Felsberg J, Stockhammer F, et al. NOA-04 randomized phase III trial of sequential radiochemotherapy of anaplastic glioma with procarbazine, lomustine, and vincristine or temozolomide. J Clin Oncol. 2009;27(35):5874-80.

52. van den Bent MJ, Dubbink HJ, Sanson M, van der Lee-Haarloo CR, Hegi M, Jeuken JW, et al. MGMT promoter methylation is prognostic but not predictive for outcome to adjuvant PCV chemotherapy in anaplastic

oligodendroglial tumors: a report from EORTC Brain Tumor Group Study 26951. *J Clin Oncol*. 2009;27(35):5881-6.

53. van den Bent MJ, Erdem-Eraslan L, Idbaih A, de Rooi J, Eilers PH, Spliet WG, et al. MGMT-STP27 methylation status as predictive marker for response to PCV in anaplastic Oligodendrogliomas and Oligoastrocytomas. A report from EORTC study 26951. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2013;19(19):5513-22.

54. Wu G, Broniscer A, McEachron TA, Lu C, Paugh BS, Becksfort J, et al. Somatic histone H3 alterations in pediatric diffuse intrinsic pontine gliomas and non-brainstem glioblastomas. *Nat Genet*. 2012;44(3):251-3.

55. Verhaak RG, Hoadley KA, Purdom E, Wang V, Qi Y, Wilkerson MD, et al. Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell*. 2010;17(1):98-110.

56. Reifenberger G, Weber RG, Riehm V, Kaulich K, Willscher E, Wirth H, et al. Molecular characterization of long-term survivors of glioblastoma using genome- and transcriptome-wide profiling. *Int J Cancer*. 2014;135(8):1822-31.

57. Vivanco I, Robins HI, Rohle D, Campos C, Grommes C, Nghiemphu PL, et al. Differential sensitivity of glioma- versus lung cancer-specific EGFR mutations to EGFR kinase inhibitors. *Cancer Discov*. 2012;2(5):458-71.

58. Wick W, Gorlia T, Bady P, Platten M, van den Bent MJ, Taphoorn MJ, et al. Phase II study of radiotherapy and temsirolimus versus radiochemotherapy with temozolomide in patients with newly diagnosed

glioblastoma without MGMT promoter hypermethylation (EORTC 26082).

Clinical cancer research : an official journal of the American Association for Cancer Research. 2016.

59. Galanis E, Buckner JC, Maurer MJ, Kreisberg JI, Ballman K, Boni J, et al. Phase II trial of temsirolimus (CCI-779) in recurrent glioblastoma multiforme: a North Central Cancer Treatment Group Study. J Clin Oncol. 2005;23(23):5294-304.

60. Wick W, Puduvalli VK, Chamberlain MC, van den Bent MJ, Carpentier AF, Cher LM, et al. Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. J Clin Oncol. 2010;28(7):1168-74.

61. Batchelor TT, Mulholland P, Neyns B, Nabors LB, Campone M, Wick A, et al. Phase III randomized trial comparing the efficacy of cediranib as monotherapy, and in combination with lomustine, versus lomustine alone in patients with recurrent glioblastoma. J Clin Oncol. 2013;31(26):3212-8.

62. Wick W, Brandes AA, Gorlia T, Bendszus M, Sahm F, Taal W, et al. Phase III trial exploring the combination of bevacizumab and lomustine in patients with first recurrence of a glioblastoma: The EORTC 26101 trial. Neuro-oncology. 2015;17:v1-v.

63. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, et al. Bevacizumab plus radiotherapy-temozolomide for newly diagnosed glioblastoma. The New England journal of medicine. 2014;370(8):709-22.

64. Gilbert MR, Dignam JJ, Armstrong TS, Wefel JS, Blumenthal DT, Vogelbaum MA, et al. A randomized trial of bevacizumab for newly diagnosed glioblastoma. *The New England journal of medicine*. 2014;370(8):699-708.

\*\* [63, 64] These studies in parallel demonstrated the lack of efficacy of anti-angiogenic therapy with bevacizumab in combination with standard chemoradiotherapy in glioblastoma patients.

65. Stupp R, Hegi ME, Gorlia T, Erridge SC, Perry J, Hong YK, et al. Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071-22072 study): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*. 2014;15(10):1100-8.

66. Sandmann T, Bourgon R, Garcia J, Li C, Cloughesy T, Chinot OL, et al. Patients With Proneural Glioblastoma May Derive Overall Survival Benefit From the Addition of Bevacizumab to First-Line Radiotherapy and Temozolomide: Retrospective Analysis of the AVAglio Trial. *J Clin Oncol*. 2015.

\* An overall survival benefit from the addition of anti-angiogenic therapy with bevacizumab to standard chemoradiotherapy was noted in glioblastomas with proneural gene expression.

67. Rohle D, Popovici-Muller J, Palaskas N, Turcan S, Grommes C, Campos C, et al. An inhibitor of mutant IDH1 delays growth and promotes differentiation of glioma cells. *Science*. 2013;340(6132):626-30.



68. Hashizume R, Andor N, Ihara Y, Lerner R, Gan H, Chen X, et al. Pharmacologic inhibition of histone demethylation as a therapy for pediatric brainstem glioma. *Nat Med*. 2014;20(12):1394-6.
69. Wolchok JD, Kluger H, Callahan MK, Postow MA, Rizvi NA, Lesokhin AM, et al. Nivolumab plus ipilimumab in advanced melanoma. *The New England journal of medicine*. 2013;369(2):122-33.
70. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *The New England journal of medicine*. 2010;363(8):711-23.
71. Ribas A, Kefford R, Marshall MA, Punt CJ, Haanen JB, Marmol M, et al. Phase III randomized clinical trial comparing tremelimumab with standard-of-care chemotherapy in patients with advanced melanoma. *J Clin Oncol*. 2013;31(5):616-22.
72. Brahmer J, Reckamp KL, Baas P, Crino L, Eberhardt WE, Poddubskaya E, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *The New England journal of medicine*. 2015;373(2):123-35.
73. Rizvi NA, Mazieres J, Planchard D, Stinchcombe TE, Dy GK, Antonia SJ, et al. Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol*. 2015;16(3):257-65.
74. Wen P, Reardon D, Phuphanich S, Aiken R, Landolfi J, Curry W, et al. A randomized double blind placebo-controlled phase 2 trial of dendritic cell

(DC) vaccine ICT-107 following standard treatment in newly diagnosed patients with GBM. *Neuro-oncology*. 2014;16:v8–v22.

\* A dendritic cell vaccination approach utilizing a combination of six peptides prolonged overall survival in this randomized, double-blinded phase II trial in newly diagnosed glioblastoma.

75. Schumacher T, Bunse L, Pusch S, Sahm F, Wiestler B, Quandt J, et al. A vaccine targeting mutant IDH1 induces antitumour immunity. *Nature*. 2014;512(7514):324-7.

76. Reardon DA, Schuster J, Tran DD, Fink KL, Nabors LB, Li G, et al. ReACT: Overall survival from a randomized phase II study of rindopepimut (CDX-110) plus bevacizumab in relapsed glioblastoma. *J Clin Oncol*. 2015;33:suppl; abstr 2009.

\* Vaccination therapy utilizing an EGFRvIII-specific peptide in combination with bevacizumab prolonged overall survival this phase II trial in recurrent glioblastoma.

77. Jena B, Dotti G, Cooper LJ. Redirecting T-cell specificity by introducing a tumor-specific chimeric antigen receptor. *Blood*. 2010;116(7):1035-44.

78. Reuss DE, Mamatjan Y, Schrimpf D, Capper D, Hovestadt V, Kratz A, et al. IDH mutant diffuse and anaplastic astrocytomas have similar age at presentation and little difference in survival: a grading problem for WHO. *Acta Neuropathol*. 2015;129(6):867-73.

\* The superiority of molecular marker diagnostics over histological tumor grade in IDH-mutant diffuse gliomas is underpinned by this study.

79. Buckner JC, Shaw EG, Pugh SL, Chakravarti A, Gilbert MR, Barger GR, et al. Radiation plus Procarbazine, CCNU, and Vincristine in Low-Grade Glioma. *The New England journal of medicine*. 2016;374(14):1344-55.

\*\* The efficacy of polychemotherapy with PCV in histological WHO grade II diffuse gliomas was defined with this phase III trial.

80. van den Bent JM, Erridge S, Vogelbaum MA, Nowak AK, Sanson M, Brandes AA, et al. Results of the interim analysis of the EORTC randomized phase III CATNON trial on concurrent and adjuvant temozolomide in anaplastic glioma without 1p/19q co-deletion: An Intergroup trial. *J Clin Oncol*. 2016;34:suppl; abstr LBA2000.

\*\* The efficacy of combined chemoradiotherapy in 1p/19q non-codeleted histological WHO grade III diffuse gliomas was defined with this phase III trial.

81. Okamoto Y, Di Patre PL, Burkhard C, Horstmann S, Jourde B, Fahey M, et al. Population-based study on incidence, survival rates, and genetic alterations of low-grade diffuse astrocytomas and oligodendrogliomas. *Acta Neuropathol*. 2004;108(1):49-56.

82. Reuss DE, Kratz A, Sahm F, Capper D, Schrimpf D, Koelsche C, et al. Adult IDH wild type astrocytomas biologically and clinically resolve into other tumor entities. *Acta Neuropathol*. 2015;130(3):407-17.

83. Lassman AB, Iwamoto FM, Cloughesy TF, Aldape KD, Rivera AL, Eichler AF, et al. International retrospective study of over 1000 adults with anaplastic oligodendroglial tumors. *Neuro-oncology*. 2011;13(6):649-59.

84. Wick W, Meisner C, Hentschel B, Platten M, Schilling A, Wiestler B, et al. Prognostic or predictive value of MGMT promoter methylation in gliomas depends on IDH1 mutation. *Neurology*. 2013;81(17):1515-22.
85. Nobusawa S, Watanabe T, Kleihues P, Ohgaki H. IDH1 mutations as molecular signature and predictive factor of secondary glioblastomas. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2009;15(19):6002-7.
86. Gramatzki D, Dehler S, Rushing EJ, Zaugg K, Hofer S, Yonekawa Y, et al. Glioblastoma in the Canton of Zurich, Switzerland revisited: 2005 to 2009. *Cancer*. 2016.
87. Ohgaki H, Kleihues P. Population-based studies on incidence, survival rates, and genetic alterations in astrocytic and oligodendroglial gliomas. *J Neuropathol Exp Neurol*. 2005;64(6):479-89.
88. Coons SW, Johnson PC, Scheithauer BW, Yates AJ, Pearl DK. Improving diagnostic accuracy and interobserver concordance in the classification and grading of primary gliomas. *Cancer*. 1997;79(7):1381-93.
89. Miller CR, Dunham CP, Scheithauer BW, Perry A. Significance of necrosis in grading of oligodendroglial neoplasms: a clinicopathologic and genetic study of newly diagnosed high-grade gliomas. *J Clin Oncol*. 2006;24(34):5419-26.